

Noncommutative Gröbner Bases for Almost Commutative Algebras*

Huishi Li

Department of Applied Mathematics
College of Information Science and Technology
Hainan University
Haikou 570228, China

Abstract. Let K be an infinite field and $K\langle X \rangle = K\langle X_1, \dots, X_n \rangle$ the free associative algebra generated by $X = \{X_1, \dots, X_n\}$ over K . It is proved that if I is a two-sided ideal of $K\langle X \rangle$ such that the K -algebra $A = K\langle X \rangle/I$ is almost commutative in the sense of [3], namely, with respect to its standard $\mathbb{N}$ -filtration FA , the associated $\mathbb{N}$ -graded algebra $G(A)$ is commutative, then I is generated by a finite Gröbner basis. Therefore, every quotient algebra of the enveloping algebra $U(\mathfrak{g})$ of a finite dimensional K -Lie algebra $\mathfrak{g}$ is, as a noncommutative algebra of the form $A = K\langle X \rangle/I$, defined by a finite Gröbner basis in $K\langle X \rangle$.

2000 Mathematics Classification Primary 16W70; Secondary 16Z05.

Key words Almost commutative algebra, Filtration, gradation, Gröbner basis

1. Introduction

Let K be an infinite field and let $K\langle X \rangle = K\langle X_1, X_2, \dots, X_n \rangle$ be the free associative K -algebra generated by $X = \{X_1, X_2, \dots, X_n\}$ over K . It is a well-known fact that even if a two-sided ideal J of $K\langle X \rangle$ is finitely generated, J does not necessarily have a finite Gröbner basis in the sense of [9]. However, it was proved in [4] that if the quotient algebra $A = K\langle X \rangle/J$ is commutative, then, after a general linear change of variables (if necessary), J has a finite Gröbner basis in $K\langle X \rangle$ (the construction of a Gröbner basis for J is mentioned in the beginning of section 3 below). This, indeed, makes another algorithmic way to study a commutative algebra via its noncommutative Gröbner presentation, and its effectiveness may be illustrated, for example, by

*Project supported by the National Natural Science Foundation of China (10571038).

the work of [1], [2], [5] and [11]. In this note, we use the result of [4] and the filtered-graded transfer of Gröbner bases [7] to show that for a two-sided ideal I of $K\langle X \rangle$, if the K -algebra $A = K\langle X \rangle/I$ is almost commutative in the sense of [3], namely, with respect to its standard $\mathbb{N}$ -filtration FA , the associated $\mathbb{N}$ -graded algebra $G(A)$ is commutative, then I is generated by a finite Gröbner basis. After reaching the main result in section 2, we discuss in section 3 how this result may be realized computationally.

Throughout the note, we fix the infinite field K , the finite set $X = \{X_1, X_2, \dots, X_n\}$ and the corresponding free K -algebra $K\langle X \rangle = K\langle X_1, X_2, \dots, X_n \rangle$.

2. The Main Result

Since the free K -algebra $K\langle X \rangle = K\langle X_1, X_2, \dots, X_n \rangle$ has its standard K -basis $\mathcal{B}$ consisting of all monomials (words) $X_{j_1}X_{j_2}\cdots X_{j_s}$, $s \in \mathbb{N}$, to be convenient, we denote monomials in $\mathcal{B}$ by $u, v, w, \dots$. Consider the $\mathbb{N}$ -gradation $K\langle X \rangle = \bigoplus_{p \in \mathbb{N}} K\langle X \rangle_p$ of $K\langle X \rangle$ with $K\langle X \rangle_p = K\text{-Span}\{u \in \mathcal{B} \mid d(u) = p\}$, where $d(u)$ stands for the degree (length) of u . If I is a two-sided ideal of $K\langle X \rangle$, then the K -algebra $A = K\langle X \rangle/I$ has the *standard $\mathbb{N}$ -filtration*

$$F_0A = K \subset F_1A \subset \cdots \subset F_pA \subset \cdots$$

where for each $p \in \mathbb{N}$,

$$F_pA = (\bigoplus_{k \leq p} K\langle X \rangle_k + I)/I,$$

which defines the associated $\mathbb{N}$ -graded K -algebra $G(A) = \bigoplus_{p \in \mathbb{N}} G(A)_p$ of A with $G(A)_p = F_pA/F_{p-1}A$. Recall from the literature [3] that A is called an *almost commutative algebra* if the associated graded K -algebra $G(A)$ of A is commutative. In [3] it was proved that *A is almost commutative if and only if A is a homomorphic image of the universal enveloping algebra $U(\mathfrak{g})$ of some finite dimensional K -Lie algebra $\mathfrak{g}$* . For instance, the n -th Weyl algebra $A_n(K)$ is an almost commutative algebra and it is a homomorphic image of the $2n+1$ -dimensional Heisenberg Lie algebra. So the class of almost commutative K -algebras consists of quotient algebras of enveloping algebras of finite dimensional K -Lie algebras. It is well-known that the study of quotient algebras of enveloping algebras (for example, the primeness, primitivity, simplicity, etc) has been very important in the finite dimensional Lie-theory. The reader is referred to ([10] Chapter 8 and Chapter 14) for details on this topic. The result and its proof given below provide a way to study the structural properties of almost commutative K -algebras algorithmically, or more precisely, to study such algebras by using both commutative and noncommutative Gröbner bases.

2.1. Theorem If the K -algebra $A = K\langle X \rangle/I$ is almost commutative, then I has a finite Gröbner basis. Equivalently, for a finite dimensional K -Lie algebra $\mathfrak{g}$, every quotient algebra of the enveloping algebra $U(\mathfrak{g})$ of $\mathfrak{g}$, viewed as a quotient of some free K -algebra, is defined by a finite Gröbner basis.

Proof For an element $f \in K\langle X \rangle$, if $f = F_p + F_{p-1} + \cdots + F_0$ with $F_i \in K\langle X \rangle_i$ and $F_p \neq 0$, then we write $d(f) = p$ for the degree of f and $\mathbf{LH}(f) = F_p$ for the leading homogeneous part of f . Consider the standard $\mathbb{N}$ -filtration FA of A and its associated $\mathbb{N}$ -graded algebra $G(A)$, as mentioned before. By ([7] Chapter III Proposition 3.1, or [8]), $G(A) \cong K\langle X \rangle / \langle \mathbf{LH}(I) \rangle$, where $\langle \mathbf{LH}(I) \rangle$ is the graded two-sided ideal of $K\langle X \rangle$ generated by $\mathbf{LH}(I) = \{\mathbf{LH}(f) \mid f \in I\}$. Since A is almost commutative, $K\langle X \rangle / \langle \mathbf{LH}(I) \rangle$ is a commutative K -algebra.

Now, let $K[\mathbf{x}] = K[x_1, x_2, \dots, x_n]$ be the commutative polynomial K -algebra in variables $x_1, x_2, \dots, x_n$ and consider the natural K -algebra homomorphism $\gamma: K\langle X \rangle \rightarrow K[\mathbf{x}]$ with $\gamma(X_i) = x_i$. Then $G(A) \cong K\langle X \rangle / \langle \mathbf{LH}(I) \rangle \cong K[\mathbf{x}] / \gamma(\langle \mathbf{LH}(I) \rangle)$ and by ([4] Theorem 2.1, Corollary 1.1), a finite Gröbner basis of $\langle \mathbf{LH}(I) \rangle$ may be obtained by using a finite Gröbner basis of $\gamma(\langle \mathbf{LH}(I) \rangle)$. Our aim is to lift the obtained Gröbner basis of $\langle \mathbf{LH}(I) \rangle$ to a finite Gröbner basis of I as described in ([7] Chapter III section 3). To this end, we need to use a graded monomial ordering. Note that γ does not change the degree (length) of monomials, that is, for any two monomials u, v in the standard K -basis $\mathcal{B}$ of $K\langle X \rangle$,

$$d(u) < d(v) \text{ if and only if } d(\gamma(u)) < d(\gamma(v)).$$

It turns out that if we fix a certain graded monomial ordering $\prec$ (for example, the graded lexicographic ordering) on $K[\mathbf{x}]$, then the *lexicographic extension* $\prec_{et}$ of $\prec$ to $K\langle X \rangle$ (in the sense of [4]), which is defined for $u, v \in \mathcal{B}$ by

$$u \prec_{et} v \text{ if } \begin{cases} \gamma(u) \prec \gamma(v) \text{ or} \\ \gamma(u) = \gamma(v) \text{ and } u \text{ is lexicographically smaller than } v, \end{cases}$$

is a graded monomial ordering. Thus, it follows from ([4] Theorem 2.1, Corollary 1.1) that $\langle \mathbf{LH}(I) \rangle$ has a finite Gröbner basis G with respect to $\prec_{et}$. Since $\gamma(\langle \mathbf{LH}(I) \rangle)$ is a graded ideal of $K[\mathbf{x}]$, by the construction of G given in ([4] Theorem 2.1, or see the description given in the beginning of section 3 below), we may assume that $G = \{G_1, G_2, \dots, G_s\}$ consists of homogeneous elements in $\langle \mathbf{LH}(I) \rangle$. Suppose for $i = 1, 2, \dots, s$,

$$G_i = \sum H_i \mathbf{LH}(f_i) T_i, \text{ where } H_i, T_i \text{ are homogeneous elements and } f_i \in I.$$

If we put $f_i = \mathbf{LH}(f_i) + f'_i$, where $d(f'_i) < d(f_i)$, then $\sum H_i f_i T_i = g_i \in I$ and

$$g_i = \sum H_i f_i T_i = \sum H_i \mathbf{LH}(f_i) T_i + \sum H_i f'_i T_i = G_i + \sum H_i f'_i T_i$$

has its leading homogeneous part $\mathbf{LH}(g_i) = G_i$, $i = 1, 2, \dots, s$. As $\prec_{et}$ is a graded monomial ordering on $K\langle X \rangle$, it follows from ([7] Chapter III Theorem 3.7) that $\mathcal{G} = \{g_1, g_2, \dots, g_s\}$ is a finite Gröbner basis of I with respect to $\prec_{et}$. This completes the proof. $\square$

3. Further Discussion

Concerning the computational realization of Theorem 2.1, we first recall from [4] how a noncommutative Gröbner basis may be constructed by using a commutative version. Let

$K\langle X \rangle = K\langle X_1, X_2, \dots, X_n \rangle$, $K[\mathbf{x}] = K[x_1, x_2, \dots, x_n]$, and $\gamma: K\langle X \rangle \rightarrow K[\mathbf{x}]$ with $\gamma(X_i) = x_i$ be as before. The *lexicographic splitting* of γ is defined as the K -linear map

$$\begin{aligned} \delta: K[\mathbf{x}] &\longrightarrow K\langle X \rangle \\ x_{i_1}x_{i_2}\cdots x_{i_r} &\mapsto X_{i_1}X_{i_2}\cdots X_{i_r} \quad i_1 \leq i_2 \leq \cdots \leq i_r \end{aligned}$$

For a monomial ideal L in $K[\mathbf{x}]$, if $m = x_{i_1}\cdots x_{i_r} \in L$ with $i_1 \leq \cdots \leq i_r$, let $\mathcal{U}_L(m)$ denote the set of all monomials $u \in K[x_{i_1+1}, \dots, x_{i_r-1}]$ such that neither $u \frac{m}{x_{i_1}}$ nor $u \frac{m}{x_{i_r}}$ lies in L . Following ([4] Theorem 2.1), if $\Gamma = \{g_1, g_2, \dots, g_s\}$ is a minimal $\prec$ -Gröbner basis for an ideal $I \subset K[\mathbf{x}]$, then a minimal $\prec_{et}$ -Gröbner basis for $J = \gamma^{-1}(I) \subset K\langle X \rangle$ consists of $\{X_jX_i - X_iX_j \mid 1 \leq i < j \leq n\}$ together with the elements $\delta(u \cdot g_i)$ for each $g_i \in \Gamma$ and each $u \in \mathcal{U}_{\langle \mathbf{LH}(I) \rangle}(\mathbf{LM}(g_i))$, where $\mathbf{LM}(g_i)$ stands for the $\prec$ -leading monomial of g_i and $\langle \mathbf{LM}(I) \rangle$ is the monomial ideal generated by $\mathbf{LM}(I) = \{\mathbf{LM}(f) \mid f \in I\}$.

In view of the characterization of an almost commutative algebra [3], to realize Theorem 2.1 computationally, we have to consider two coherent cases separately. In what follows, notations used in section 2 are maintained.

I. $A = K\langle X \rangle / I$, $I = \langle S \rangle$ and $G(A)$ is commutative.

Since $G(A)$ is commutative, it follows from the proof of Theorem 2.1 that we have

$$G(A) \cong K\langle X \rangle / \langle \mathbf{LH}(I) \rangle \cong K[\mathbf{x}] / \gamma(\langle \mathbf{LH}(I) \rangle).$$

Again by the proof of Theorem 2.1, in order to obtain a finite Gröbner basis of I algorithmically, we need to have a finite Gröbner basis of $\gamma(\langle \mathbf{LH}(I) \rangle)$ so that we may use it to construct a Gröbner basis for $\langle \mathbf{LH}(I) \rangle$ as remarked above. While this implies that we need first to know a generating set of $\gamma(\langle \mathbf{LH}(I) \rangle)$. But the fact is that even if the generating set of I is finite, say $S = \{f_1, f_2, \dots, f_s\}$, the equality $\langle \mathbf{LH}(I) \rangle = \langle \mathbf{LH}(f_1), \mathbf{LH}(f_2), \dots, \mathbf{LH}(f_s) \rangle$ is not necessarily true (see [7] Chapter III). In other words, if we cannot find a generating set of $\gamma(\langle \mathbf{LH}(I) \rangle)$ effectively, Theorem 2.1 is only theoretical.

As an example, let us point out that if we know some (finite or infinite) Gröbner basis $\mathcal{G} = \{g_i \mid i \in \Omega\}$ of I and $\mathcal{G}$ contains all commutators $X_iX_j - X_jX_i$, $1 \leq i < j \leq n$, then $\langle \mathbf{LH}(I) \rangle = \langle \mathbf{LH}(g_i) \mid i \in \Omega \rangle$ by ([7] Chapter III).

II. $A = U(\mathfrak{g})/I$ and $I = \langle S \rangle$.

Suppose $U(\mathfrak{g}) = K\langle X \rangle / J$ with $J = \langle [X_i, X_j] - X_iX_j - X_jX_i \mid 1 \leq i < j \leq n \rangle$, where $[X_i, X_j]$ is given by the Lie-brackets of generators of the finite dimensional K -Lie algebra $\mathfrak{g}$. Let $\bar{I}$ be the two-sided ideal of $K\langle X \rangle$ such that $I = \bar{I}/J$. Then $A = U(\mathfrak{g})/I \cong K\langle X \rangle / \bar{I}$. By the foregoing discussion, if we can find a finite Gröbner basis of $\gamma(\langle \mathbf{LH}(\bar{I}) \rangle)$, then a finite Gröbner basis for $\bar{I}$ may be constructed.

Note that the standard filtration $FU(\mathfrak{g})$ induces the standard filtration FA . Consider the filtration FI on I induced by $FU(\mathfrak{g})$ and its associated graded ideal $G(I)$ in $G(A)$. Then it is

well known that

$$G(A) = G(U(\mathfrak{g})/I) \cong G(U(\mathfrak{g}))/G(I) \cong K[\mathbf{x}]/\overline{G(I)},$$

where $\overline{G(I)}$ stands for the counterpart of $G(I)$ in $K[\mathbf{x}]$. Thus, since both graded algebra epimorphisms

$$\varphi : k[\mathbf{x}] \longrightarrow K[\mathbf{x}]/\overline{G(I)} \cong G(A)$$

$$\psi : k[\mathbf{x}] \longrightarrow K[\mathbf{x}]/\gamma(\langle \mathbf{LH}(\bar{I}) \rangle) \cong G(A)$$

agree on the generators $x_1, x_2, \dots, x_n$, we have $\overline{G(I)} = \gamma(\langle \mathbf{LH}(\bar{I}) \rangle)$. This makes the chance for us to have a finite Gröbner basis of $\gamma(\langle \mathbf{LH}(\bar{I}) \rangle)$ by using the generating set S of I . To see this, first recall that $U(\mathfrak{g})$ is a solvable polynomial algebra in the sense of [6]. Hence, starting with S , a noncommutative version of Buchberger Algorithm produces a finite Gröbner basis $\mathcal{G} = \{g_1, g_2, \dots, g_m\}$ for I . Furthermore, if the monomial ordering used in producing $\mathcal{G}$ is a graded monomial ordering, then it follows from ([7] Chapter IV Theorem 2.1) that $\sigma(\mathcal{G}) = \{\sigma(g_1), \sigma(g_2), \dots, \sigma(g_m)\}$ is a Gröbner basis for $G(I)$ with respect to the same type of graded monomial ordering, where for each g_i , if $g_i \in F_p I - F_{p-1} I$, then $\sigma(g_i)$ is the homogeneous element in $G(I)_p$ represented by g_i . Finally, by passing to $\overline{G(I)}$ in $K[\mathbf{x}]$, we obtain a finite Gröbner basis $\{\bar{g}_1, \bar{g}_2, \dots, \bar{g}_m\}$ of $\gamma(\langle \mathbf{LH}(\bar{I}) \rangle)$. Note that $U(\mathfrak{g}) = K\langle X \rangle/J$, $I = \bar{I}/J$. Hence the preimage of $\mathcal{G}$ is contained in $\bar{I}$. Thus, by the definition of $\delta(u \cdot \bar{g}_i)$ described in the beginning of this section, the last step of the proof of Theorem 2.1 can be realized to give a Gröbner basis for $\bar{I}$.

In this case, the good news is that nowadays there has been the well-developed computer algebra system SINGULAR: PLURAL (<http://www.singular.uni-kl.de/index.html>) which provides a programme named **twostd** for computing a two-sided Gröbner basis of a two-sided ideal in a solvable polynomial algebra.

We end this note by employing an easy example to illustrate the procedure demonstrated in part I and II above. Let $U(sl_2)$ be the enveloping algebra of the 3-dimensional K -Lie algebra $sl_2 = Ke \oplus Kf \oplus Kh$ subject to the relations $[e, f] = h$, $[h, e] = 2e$, $[h, f] = -2f$, that is, $U(sl_2) = K[e, f, h] \cong K\langle X, Y, Z \rangle/P$ with $X \mapsto e$, $Y \mapsto f$ and $Z \mapsto h$, where

$$P = \langle YX - XY + Z, ZX - XZ - 2X, ZY - YZ + 2Y \rangle.$$

Consider the two-sided ideal $I = \langle e^3, f^3, h^3 - 4h \rangle$ of $U(\mathfrak{g})$, then $U(\mathfrak{g})/I \cong K\langle X, Y, Z \rangle/\bar{I}$, where

$$\bar{I} = \langle YX - XY + Z, ZX - XZ - 2X, ZY - YZ + 2Y, X^3, Y^3, Z^3 - 4Z \rangle.$$

Let $\prec$ be the graded reverse lexicographic ordering on $U(\mathfrak{g})$ defined by $e \prec f \prec h$. Then **twostd** produces a two-sided Gröbner basis

$$\mathcal{G} = \left\{ \begin{array}{l} e^3, f^3, h^3 - 4h, eh^2 + 2eh, \\ fh^2 - 2fh, 2efh - h^2 - 2h, \\ e^2f - eh - 2e, ef^2 - fh, e^2h + 2e^2, f^2h - 2f^2 \end{array} \right\}$$

for I (Singular Manual A.6.1). Hence

$$\sigma(\mathcal{G}) = \left\{ \begin{array}{l} \sigma(e)^3, \sigma(f)^3, \sigma(h)^3, \sigma(e)\sigma(h)^2, \sigma(f)\sigma(h)^2, \\ 2\sigma(e)\sigma(f)\sigma(h), \sigma(e)^2\sigma(f), \sigma(e)\sigma(f)^2, \sigma(e)^2\sigma(h), \sigma(f)^2\sigma(h) \end{array} \right\}$$

is a Gröbner basis of $G(I)$. If we use the graded reverse lexicographic ordering $x \prec y \prec z$ on the polynomial K -algebra $K[x, y, z]$, then previous part II yields a Gröbner basis

$$\{x^3, y^3, z^3, xz^2, yz^2, 2xyz, x^2y, xy^2, x^2z, y^2z\}$$

for $\gamma(\langle \mathbf{LH}(\bar{I}) \rangle)$. By the construction of $\delta(u \cdot g)$ described in the beginning of this section, it may be checked directly that $\langle \mathbf{LH}(\bar{I}) \rangle$ has a Gröbner basis

$$\left\{ \begin{array}{l} YX - XY, ZY - YZ, ZX - XZ, X^3, Y^3, \\ Z^3, XZ^2, YZ^2, 2XYZ, X^2Y, XY^2, X^2Z, Y^2Z \end{array} \right\}.$$

Note that $\bar{I}$ contains the preimage of $\mathcal{G}$. It follows from the last step of the proof of Theorem 2.1 that $\bar{I}$ has the Gröbner basis

$$\left\{ \begin{array}{l} YX - XY, ZY - YZ, ZX - XZ, \\ X^3, Y^3, Z^3 - 4Z, XZ^2 + 2XZ, \\ YZ^2 - 2YZ, 2XYZ - Z^2 - 2Z, \\ X^2Y - XZ - 2X, XY^2 - YZ, X^2Z + 2X^2, Y^2Z - 2Y^2 \end{array} \right\}$$

with respect to $X \prec_{et} Y \prec_{et} Z$.

References

- [1] D. J. Anick, On the homology of associative algebras, *Trans. Amer. Math. Soc.*, Vol. 296, 2(1986), 641–659.
- [2] D. Anick and G.-C. Rota, Higher-order syzygies for the bracket algebra and for the ring of coordinates of the Grassmannian, *Proc. Nat. Acad. Sci. U.S.A.*, 88(1991), 8087C8090.
- [3] M. Duflo, Certaines algèbres de type finissent des algèbres de Jacobson, *J. Algebra*, 27(1973), 358–365.
- [4] D. Eisenbud, I. Peeva and B. Sturmfels, Non-commutative Gröbner bases for commutative algebras, *Proc. Amer. Math. Soc.*, 126(1998), 687–691.
- [5] E. Green and R.Q. Huang, Projective resolutions of straightening closed algebras generated by minors, *Adv. in Math.*, 110(1995), 314C333.
- [6] A. Kandri-Rody and V. Weispfenning, Non-commutative Gröbner bases in algebras of solvable type, *J. Symbolic Comput.*, 9(1990), 1–26.
- [7] Huishi Li, *Noncommutative Gröbner Bases and Filtered-Graded Transfer*, LNM, 1795, Springer-Verlag, 2002.
- [8] Huishi Li, The general PBW property, *Algebra Colloquium*, in press.
- [9] T. Mora, An introduction to commutative and noncommutative Gröbner bases, *Theoretical Computer Science*, 134(1994), 131–173.

- [10] J.C. McConnell and J.C. Robson, *Noncommutative Noetherian Rings*, John Wiley & Sons, 1987.
- [11] I. Peeva, V. Reiner and B. Sturmfels, How to shell a monoid, *Math. Ann.*, 310(1998), 379C393.